

Development of Antibodies That Show Antigen-dependent Fluorescence Enhancement

○Keisuke Fukunaga¹, Takayoshi Watanabe¹, Dian Novitasari¹, Ryoji Abe², Hiroyuki Ohashi², and Takahiro Hohsaka^{1,*}

¹ School of Materials Science, Japan Advanced Institute of Science and Technology (JAIST)
Nomi, Ishikawa 923-1292, Japan. *E-mail: hohsaka@jaist.ac.jp ² USHIO Inc.




Abstract

Protein-based fluorescent biosensor is an undoubtedly powerful tool for detection and visualization of target biomolecules. Recently, it was reported that single-chain variable fragment (scFv) antibody-based fluorescent biosensor, namely Quenchbody showed a fluorescence intensity change in an antigen concentration-dependent manner. Because structurally conserved protein (*i.e.*, antibody) was employed as the scaffold of biosensor, measurements of various antigens were achieved by preparing fluorescent amino acid-incorporated scFv. However, time-consuming gene manipulation and usage of less productive cell-free translation system is required for genetic incorporation of fluorescent amino acid into scFv. Thus, alternative approach for preparation of antibody-based fluorescent biosensor is highly demanded.


We here report a novel design and synthesis of the antibody-based fluorescent biosensor. Previous study proved that fluorescent amino acid at the N-terminal side of scFv was quenched by conserved tryptophans, but the binding of its antigen triggered a structural rearrangement of scFv followed by a cancellation of the fluorescence quenching. Accordingly, it is considered that N-terminal-selective fluorescent labeling will be useful for preparation of antibody-based biosensor. As a proof of concept, immunoglobulin (IgG) monoclonal antibody-based fluorescent biosensor was semi-synthesized by reductive alkylation of N-terminal α -amino groups of IgG with dye-aldehyde in a weak acidic buffer. The N-terminal-selective conjugation of fluorescent dye was evaluated by tandem mass spectrometric analysis of peptide fragments. Also, antigen-dependent fluorescence intensity change of the labeled IgG was confirmed by fluorescence spectral measurement. Fluorescent detection of several peptides and biomolecules was successfully achieved by using the IgG-based biosensor. Because a variety of different IgG are commercially available, wide applications of the IgG-based biosensor will be developed.

Research background


1) Incorporation of NAA into N-terminus of scFv by expanded genetic code system



2) Fluorescence detection of controlled drugs using the fluorescent-labeled scFv



3) TAMRA on the N-terminal domain is quenched by intrinsic conserved Trp residues



- Rapid
- Quantitative
- In solution (immobilization & wash-free)
- Small molecules can be detected.
- Various antigens are available.




Reference

1) Ryoji Abe *et al.*, J. Am. Chem. Soc. 133, 17386 (2011)


Results

1) N-terminal α -amino group-selective fluorescent-labeling




pK_a values
• α -amino group: 7.7 ± 0.5
• ϵ -amino group of lysine: 10.5 ± 1.1
cited from J. Biol. Chem. 284, 13285 (2009)

2) Synthesis and characterization of TAMRA-labeled anti-FLAG IgG




3) Optimization of fluorescent dye and alkyl linker length




Positioning of this study

Schematic drawing of synthesis of antibody-based biosensors: strategies for incorporation of fluorescent dye



4) MS/MS analysis of the trypsinized IgG (anti-c-Myc 9E10 monoclonal)



5) TAMRA-labeled IgG antibodies might expand various applications

